

Document Number: Version Number: Effective date: Written by: Approved by: QF0244_Kit_CoA_Cover 0 2016-05-16 QC Manager Production Manager

Kit CoA Cover Page

| KAPA HiFi HS Uracil+ RM (0.25 ml SMP) | | | | |
|---------------------------------------|-------------|-------------|-----------------|--|
| Kit Code | Part Number | Lot Number | Kit Expiry Date | |
| KK2800 | 07959044001 | 004373-14-1 | 2018-04-10 | |

| Component Code | Component Description | Component Lot Number |
|----------------|---------------------------------|----------------------|
| KM2800 | 2x HiFi HS Uracil+ MM (0.25 ml) | 00063099 |

CoA's are not issued for complete kits, but for the individual component lots from which kits are assembled. CoA's for all component lots listed are attached.

| Generated By | Date | |
|----------------------------------|------------|--|
| Danielle Da Silva (QC Scientist) | 2016-10-03 | |



Certificate of Analysis

| PRODUCT DETAILS | | | | |
|--|------------------------------|---------|--|--|
| Product name | 2X HiFi HS Uracil + ReadyMix | | | |
| Code & Pack size | KM2800 | 0.25 ml | | |
| Lot number | 63099 | | | |
| Code and lot number of bulk corresponding solution | BM0030 | 61880 | | |

| QUALITY CONTROL PARAMETERS | | | |
|----------------------------|---|--------|--|
| Parameter | Specification | | |
| Purity | The enzyme contained in this product is extensively purified through the use of multiple chromatography steps. The final formulation contains <2% contaminating protein, as determined in an Agilent Protein 230 Assay. The nucleotides contained in this product are >98% pure, as determined by HPLC analysis. | Passed | |
| Functional assay | A single, distinct band visible by agarose gel electrophoresis/ ethidium bromide staining, following amplification of a 599 bp DNA fragment from a dilution series of 10 ng – 100 pg human genomic DNA under standard reaction conditions. | Passed | |
| Endonuclease activity | No detectable nicking or linearization of plasmid DNA after incubation for 8 hours at 37 °C, as assessed by agarose gel electrophoresis/ethidium bromide staining. | Passed | |
| Exonuclease activity | No detectable degradation of lambda DNA after incubation for 8 hours at 37 $^{\circ}$ C, as assessed by agarose gel electrophoresis/ ethidium bromide staining. | Passed | |
| DNA contamination | A standard reaction with no template contains <50 fg/µl bacterial genomic DNA (<i>E. coli</i> and related strains; as assessed by amplification of a 411 bp 16S rRNA fragment using a multicopy primer set in a 35-cycle reaction) and <0.5 pg/µl human genomic DNA (as assessed by amplification of a 290 bp b-actin fragment using a multicopy primer set in a 35-cycle reaction). | Passed | |

Generated by Toni Hartel (QC Scientist)

2015-02-13